

QCD Thermodynamics with Improved Actions

F. Karsch * with B. Beinlich, J. Engels, R. Joswig, E. Laermann, A. Peikert and B. Petersson

Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany

The thermodynamics of the SU(3) gauge theory has been analyzed with tree level and tadpole improved Symanzik actions. A comparison with the continuum extrapolated results for the standard Wilson action shows that improved actions lead to a drastic reduction of finite cut-off effects already on lattices with temporal extent $N_\tau = 4$. Results for the pressure, the critical temperature, surface tension and latent heat are presented. First results for the thermodynamics of four-flavour QCD with an improved staggered action are also presented. They indicate similarly large improvement factors for bulk thermodynamics.

1. Why improved actions are expected to improve the QCD thermodynamics

Although the plasma phase of QCD does in many respects show strong non-perturbative properties (screening masses, poor convergence of perturbation theory,...), bulk thermodynamic observables like energy density and pressure do approach the non-interacting ideal gas limit at high temperatures (T) where they are expected to receive large contributions from high momentum modes, $\langle \text{avg.momentum} \rangle \sim T$. In lattice calculations T is determined through the temporal extent N_τ of the lattice and the cut-off a^{-1} , *i.e.* $T \equiv 1/N_\tau a$. In the ideal gas limit the relevant momenta are therefore of the order of the cut-off, where lattice and continuum dispersion relations differ strongly from each other. Indeed this is the origin of the well-known discrepancy between the energy density of an ideal gas calculated on a finite lattice ($\epsilon(N_\tau)$) and in the continuum (ϵ_{SB}),

$$\frac{\epsilon(N_\tau)}{\epsilon_{\text{SB}}} = \begin{cases} 1 + \frac{30}{63} \cdot \frac{\pi^2}{N_\tau^2} + \mathcal{O}(N_\tau^{-4}) & , \text{ Wilson} \\ 1 + c_I \cdot \frac{\pi^4}{N_\tau^4} + \mathcal{O}(N_\tau^{-6}) & , \text{ Symanzik} \end{cases}$$

The standard Wilson action leads to nearly 50% corrections on a lattice with temporal extent $N_\tau = 4$ while the cut-off dependence is drastically reduced in the case of Symanzik improved actions which eliminate the leading $\mathcal{O}(N_\tau^{-2}) \equiv \mathcal{O}((aT)^2)$

*The work has been supported by the DFG under grants Pe 340/3-3 and Pe 340/6-1.

corrections. In the case of the (1,2)-Symanzik action this is less than 2% for $N_\tau = 4$, *i.e.* $c_I = 0.044$. This clearly demonstrates the significance of improved actions in the ideal gas limit ($T \rightarrow \infty$).

In a first exploratory analysis we have shown that the improvement found analytically in the ideal gas limit also persists at finite temperature and does seem to be important even close to T_c [1]. Here we will present further evidence that tree level and tadpole improved actions do lead to a significant reduction of finite cut-off effects even at T_c [2-4].

2. SU(3) Thermodynamics

We have analyzed the thermodynamics of the SU(3) gauge theory using $\mathcal{O}(a^2)$ improved tree level and tadpole improved actions, which have been constructed by adding to the Wilson plaquette action, $S^{(1,1)}$, an appropriately weighted contribution from planar (1,2) or (2,2) loops,

$$S^{(1,2)} = \sum_{x,\nu>\mu} \frac{5}{3} \left(1 - \frac{1}{N} \text{Re Tr } \square_{\mu\nu}(x) \right) \quad (1)$$

$$- \frac{1}{6u_0^2} \left(1 - \frac{1}{2N} \text{Re Tr } \left(\square_{\mu\nu}(x) + \square_{\nu\mu}(x) \right) \right)$$

$$S^{(2,2)} = \sum_{x,\nu>\mu} \frac{4}{3} \left(1 - \frac{1}{N} \text{Re Tr } \square_{\mu\nu}(x) \right) \quad (2)$$

$$- \frac{1}{48} \left(1 - \frac{1}{N} \text{Re Tr } \square_{\mu\nu}(x) \right)$$

Table 1

Critical temperature in units of $\sqrt{\sigma}$ on lattices with temporal extent $N_\tau = 4$. Infinite volume extrapolations of β_c for the Symanzik actions are based on lattices with $N_\sigma = 16, 24$ and 32 .

action	N_σ	β_c	$T_c/\sqrt{\sigma}$
standard Wilson	∞	5.69254 (24)	0.5983 (30)
(2,2) Symanzik (tree level)	24	4.3995 (2)	0.624 (4)
(1,2) Symanzik (tree level)	∞	4.07297 (28)	0.631 (3)
(1,2) Symanzik (tadpole)	∞	4.35228 (39)	0.635 (3)

where the factor u_0 appearing in the definition of the (1,2)-Symanzik action denotes the tadpole improvement factor defined in terms of the self-consistently determined plaquette expectation value [7].

2.1. Bulk thermodynamics

Bulk thermodynamic quantities like the energy density (ϵ) or pressure (p) calculated with the Wilson action on lattices of size $N_\sigma^3 \times N_\tau$ with $N_\tau = 4, 6$ and 8 [5] show a strong cut-off dependence in the plasma phase. A first analysis with the (2,2)-Symanzik action [1] has shown that this cut-off dependence gets drastically reduced already on a $N_\tau = 4$ lattice. The work presented in Ref. [1] has been extended in several ways. On lattices with temporal extent $N_\tau = 4$ we have analyzed now also the tree-level improved (1,2)-Symanzik action, which shows even smaller cut-off distortions in the high temperature limit than the (2,2) action. In addition we have studied the influence of tadpole improvement in the case of the (1,2)-Symanzik action. In all cases we have determined the temperature scale non-perturbatively through the calculation of the string tension on symmetric (N_σ^4) lattices, $T/T_c \equiv \sqrt{\sigma}a(\beta_c)/\sqrt{\sigma}a(\beta)$. From this calculation we also obtain T_c in units of $\sqrt{\sigma}$ for the different actions. These are given in Table 1 together with results for the Wilson action. In that case $T_c/\sqrt{\sigma}$ has also been calculated on lattices with N_τ varying between 4 and 12 and extrapolated to the continuum limit. This gave $T_c/\sqrt{\sigma} = 0.629 \pm 0.003$ [5]. On $N_\tau = 4$ lattices the results obtained with improved actions are thus much closer to this extrapolated value than those obtained with the Wilson action.

The pressure has been calculated by integrating differences of the action densities obtained on 16^4 and $16^3 \times 4$ lattices (in the case of the (2,2) action we used 24^4 and $24^3 \times 4$ lattices) following the standard procedures [1,5]. Results are shown in Figure 1.




Figure 1. Pressure of the SU(3) gauge theory on lattices with temporal extent $N_\tau = 4$. The solid line shows the continuum extrapolation obtained from the standard Wilson action. The dots give results from a calculation with a perfect action on a $12^3 \times 3$ lattice [6]. The arrow indicates the continuum ideal gas value.

2.2. Surface tension and latent heat

The success of improved actions for the calculation of bulk thermodynamics even at temperatures close to T_c naturally leads to the question whether these actions also do lead to an improvement at T_c . This is, of course, a highly non-perturbative regime. However, observables like the latent heat ($\Delta\epsilon$) and the surface tension (σ_I), which characterize the discontinuities at the first

order deconfinement phase transition in a $SU(3)$ gauge theory, do depend on properties of the low as well as the high temperature phase. As the latter is largely controlled by high momentum modes it may be expected that some improvement does result even from tree level improved actions.

At the $SU(3)$ deconfinement transition $\Delta\epsilon$ and σ_I have been studied on lattices up to temporal extent $N_\tau = 6$ [8,9]. A strong cut-off dependence has been found when comparing calculations for $N_\tau = 4$ and 6. For this reason an extrapolation to the continuum limit has so far not been possible for these observables.

We have extracted σ_I [3] from the probability distribution of the absolute value of the Polyakov loop following the analysis presented in Ref. [9]. The probability distribution at the minimum is proportional to

$$P(|L|) \sim \exp(-[f_1 V_1 + f_2 V_2 + 2\sigma_I A]/T) \quad (3)$$

where f_i denotes the free energy in the phase i , and V_i is the volume occupied by that phase and A denotes the interface area of the finite system. From the depth of the minimum one thus can extract the surface tension. The distribution functions for the tadpole improved actions for three different lattice sizes are shown in Figure 2.




Figure 2. Polyakov loop distributions for the tadpole improved (1,2) Symanzik action.

In Table 2 we give results for σ_I on the largest lattices considered. Clearly the surface tension

extracted from simulations with improved actions on lattices with temporal extent $N_\tau = 4$ are substantially smaller than corresponding results for the Wilson action. In fact, they are compatible with the $N_\tau = 6$ results for the Wilson action.

The latent heat is calculated from the discontinuity in $(\epsilon - 3P)$. This in turn is obtained from the discontinuity in the various Wilson loops entering the definition of the improved actions,

$$\frac{\Delta\epsilon}{T_c^4} = \frac{1}{6} \left(\frac{N_\tau}{N_\sigma} \right)^3 \left(a \frac{d\beta}{da} \right) \left(\langle \tilde{S} \rangle_+ - \langle \tilde{S} \rangle_- \right), \quad (4)$$

with $\tilde{S} \equiv S - dS/d\beta$.

The difference of action expectation values at β_c is obtained by calculating these in the two co-existing phases at β_c . In the time histories of the Polyakov loop values we have cut out the transition regions in order to classify configurations belonging to either of the two phases [8].

In order to extract the latent heat one does still need the β -function entering the definition of $\Delta\epsilon/T_c^4$ in Eq. 4. The necessary relation $a(\beta)$ has been obtained from a calculation of $\sqrt{\sigma}a$ (improved actions) or a determination of $T_c a$ (Wilson action). Results for $\Delta\epsilon/T_c^4$ are summarized in Table 2. Here we also give the result obtained with the 1-loop β -function.

3. Four-flavour QCD with an improved staggered action

In the fermionic sector of QCD the influence of a finite cut-off on bulk thermodynamic observables is known to be even larger than in the pure gauge sector. For instance, in the staggered formulation the energy density of an ideal fermi gas differs by more than 70% from the continuum value on a lattice with temporal extent $N_\tau = 4$ and approaches the continuum value only very slowly with increasing N_τ . This cut-off dependence can drastically be reduced with an $O(a^2)$ improved staggered action. We use an improved action, $S^I[U] = S^{(1,2)} + \bar{\psi} M \psi$, where a higher order difference scheme (one-link and three-link terms), is used to improve the fermionic part [10]. The improved fermion matrix reads

$$M[U]_{ij} = m\delta_{i,j} + \eta_i \left(\frac{9}{16} A[U]_{ij} - \frac{1}{48} B[U]_{ij} \right)$$

Table 2

Surface tension and latent heat for the three improved actions and the Wilson action. Results for the Wilson action are based on data from [9] using the non-perturbative β -function calculated in [5].

action	V_σ	N_τ	σ_I/T_c^3	$\Delta\epsilon/T_c^4$	$\left(\Delta\epsilon/T_c^4\right)_{\text{pert}}$
standard Wilson	$24^2 \times 36$	4	0.0300 (16)	2.27 (5)	4.06 (8)
	$36^2 \times 48$	6	0.0164 (26)	1.53 (4)	2.39 (6)
(1,2) Symanzik (tree level)	32^3	4	0.0116 (23)	1.57 (12)	2.28 (8)
(1,2) Symanzik (tadpole)	32^3	4	0.0125 (17)	1.40 (9)	1.94 (8)

with

$$A[U]_{ij} = \sum_{\mu} \left(U_{i,\mu} \delta_{i,j-\hat{\mu}} - U_{i-\hat{\mu},\mu}^\dagger \delta_{i,j+\hat{\mu}} \right)$$

$$B[U]_{ij} = \sum_{\mu} \left(U_{i,\mu} U_{i+\hat{\mu},\mu} U_{i+2\hat{\mu},\mu} \delta_{i,j-3\hat{\mu}} - \right.$$

$$\left. - U_{i-\hat{\mu},\mu}^\dagger U_{i-2\hat{\mu},\mu}^\dagger U_{i-3\hat{\mu},\mu}^\dagger \delta_{i,j+3\hat{\mu}} \right)$$

With this action the overall cut-off distortion of the ideal gas limit on a $16^3 \times 4$ lattices reduces to about 20%. We have performed simulations for two quark masses, $ma = 0.05$ and 0.1 [4]. Like in the pure gauge sector the improvement is visible already close to T_c . For instance, the energy density stays close to the ideal gas limit. Although we observe an overshooting of the ideal gas limit close to T_c for the non-zero quark masses considered by us this feature does seem to disappear in the "chiral limit" [4].




Figure 3. Energy density of four-flavour QCD on a $16^3 \times 4$ lattice. The lower set of curves shows an extrapolation to the "chiral limit".

4. Conclusions

Thermodynamic observables of $SU(3)$ gauge theory and QCD studied with improved gauge and fermion actions show a drastic reduction of the cut-off dependence in the high temperature limit as well as at T_c . The major improvement effect is already obtained with tree level improved actions. Tadpole improvement plays a minor role even close to T_c .

REFERENCES

1. B. Beinlich, F. Karsch and E. Laermann, Nucl. Phys. B462 (1996) 415.
2. B. Beinlich, F. Karsch, E. Laermann and A. Peikert, *String tension and Thermodynamics from Tree Level and Tadpole Improved Actions*, in preparation.
3. B. Beinlich, F. Karsch and A. Peikert, *SU(3) Latent Heat and Surface Tension from Tree Level and Tadpole Improved Actions*, BI-TP 96/24.
4. J. Engels, R. Joswig, F. Karsch, E. Laermann, and B. Petersson, *Thermodynamics of four-flavour QCD with an improved staggered action*, in preparation.
5. G. Boyd *et al.*, Phys. Rev. Lett. 75 (1995) 4169 and Nucl. Phys. B469 (1996) 419.
6. A. Papa, *SU(3) Thermodynamics on Small Lattices*, BUTP-96/13.
7. G.P. Lepage and P.B. Mackenzie, Phys. Rev. D48 (1993) 2250.
8. Y. Iwasaki *et al.*, Phys. Rev. D46 (1992) 4657.
9. Y. Iwasaki *et al.*, Phys. Rev. D49 (1994) 3540.
10. S. Naik, Nucl. Phys. B316 (1989) 238.